

## Kerry-Basic

### Benefit:

1. Improved soil structure.
2. Strong root formation, resistance to drought.
3. Increased yield.
4. Eliminates trace element deficiency.
5. Resistance to disease.
6. For ornamentals - Stronger colour, longer lasting blooms.

## 佳丽加基素

### 益处:

1. 改良土壤构造。
2. 加强根部构造, 抵抗干旱季节。
3. 增加产量。
4. 释放微量元素的不足。
5. 抵抗病害侵袭。
6. 使观赏植物颜色鲜艳, 开花更持久。



# 佳麗基素

# Kerry Basic

## Natural Seaweed Extract

纯海藻有机养素

Net Weight : 4 liter

Tarikh Mengilang:

Seaweed	10% w/v
Protein	1% w/v
Carbohydrate	10% w/v
40 Trace Elements	Al, Co, I, Mo, Na, Ni, Si, etc
Growth Stimulants	Betaines, Auxin, Cytokinins, Gibberelin.



Imported & Distributed by :

**Mycrop Sdn Bhd 国农有限公司** (507146-W)

36, Lalan Perindustrian Silibin 2,  
Off Jalan Jelapang, 30200 Ipoh, Perak.

Tel: 05-5285281 Fax: 05-5284281

FOLIAR APPLICATION (ml/4.5 liter) 施用方法 (毫升/4.5公升)

TIMING CROP 作物	First 第一 5-10 ml/毫升	Second 第二 5-10 ml/毫升	Third 第三 7.5-10 ml/毫升	Optional 施用次数 7.5-10 ml/毫升
<b>Vegetables</b> 蔬菜 Cabbage, Cauliflower, Brassica, Onion, Lettuce. 包菜, 花椰菜, 十字花科, 葱, 生菜	At 4-6 leaf stage. 4-6 片叶时	10 days later. 10 天後	At head initiation. 由头灌溉	—
<b>Fruit Vegetables</b> 果菜 Tomato, Peppers. 番茄, 甜椒	At 15-20cm growth. 生长至15-20 公分时	At first bloom. 开花初期	At first fruit set. 结果初期	Every 14 days. 每隔14 天
Beans, Peas. 豆类, 豌豆	At 4-6 leaf stage. 4-6 片叶时	At first bloom. 开花初期	At first pods. 结豆荚初期	—
Watermelon, Muskmelon. 西瓜, 蜜瓜	At 15-20cm growth. 生长至15-20 公分时	At first bloom.	At first fruit set. 结果初期	Within 48 hours of harvest. 选摘后48小时内
Cucumber 黄瓜	At 4 leaf stage. 4 片叶时	At first bloom.	7-14 days later. 7-14 天后	
<b>Fruits Tree</b> 果树 Mango, Pomelo, Guava etc.. 芒果, 柚子, 番石榴等等。	Flowering bud formation. 花蕾形成时	Flowering 开花时期	After 14 days. 14 天后	—
Paddy 稻谷	2 weeks after sowing 播种後2星期	Flowering 开花时期	After 14 days. 14 天后	—